



FEED

Performance Sheep with Deccox

Pelleted Sheep Feed (Medicated)

For the prevention of coccidiosis in young sheep caused by *Eimeria bakuensis*, *Eimeria crandalis*, *Eimeria ovinoidalis* and *Eimeria parva*. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

Feeding Directions:

This feed is formulated for sheep and should be fed at a rate of 2% of body weight per head per day.

Feed Type C feed or milk replacer at a rate to provide 22.7 mg per 100 lb (0.5 mg/kg) bodyweight. Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid type B feed.

Caution:

Do not use in feeds containing bentonite.

Always provide a source of fresh clean water, and free choice minerals. When making changes in an

animal's diet, allow a 7 to 10 day transition period. Provide ample feeder space for livestock. Do not feed wet, moldy, or insect infested feed.

Warning:

Do not feed to sheep producing milk for food.

Ingredients:


Processed Grain By-Products, Grain Products, Plant Protein Products, Forage Products, Roughage Products, Animal Protein Products, Calcium Carbonate, Potassium Sulfate, Magnesium Sulfate, Ammonium Chloride, Salt, Expand, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement and Saccharomyces Yeast Culture

Active Drug Ingredient:

Decoquinatate 27.2 g/ton

Guaranteed Analysis:

Crude Protein (min).....	18.00%
Crude Fat (min).....	3.00%
Crude Fiber (max).....	12.00%
Calcium (min).....	0.50%
Calcium (max).....	1.00%
Phosphorus (min)	0.40%
Vitamin A (min).....	18,500 IU/lb
Vitamin D (min)	2,000 IU/lb
Vitamin E (min).....	56 IU/lb



Visit your local Alabama Farmers Cooperative to learn more or visit afcfeed.com.