



Goat Feed

16% Goat with Deccox

Pelleted Goat Feed (Medicated)

For the prevention of coccidiosis in young goats caused by Eimeria christenseni and Eimeria ninakohlyakimovae. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

Feeding Directions:

This feed is to be provided for goats. Feed at the rate of 1 to 2% of body weight per head per day along with free choice hay or other forages.

Feed Type C milk replacer at a rate to provide 22.7 mg per 100 lb (0.5 mg/kg) bodyweight. Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid type B feed.

Caution:

Do not use in feeds containing bentonite.

Always provide a source of fresh clean water, and free choice minerals. When making changes in an

animal's diet, allow a 7 to 10-day transition period. Provide ample feeder space for livestock. Do not feed wet, moldy, or insect infested feed.

Warning:

Do not feed to goats producing milk for food.

Ingredients:

Processed Grain By-Products, Grain Products, Roughage Products, Plant Protein Products, Calcium Carbonate, Salt, Ammonium Chloride, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, Sodium Selenite, Ferrous Sulfate, Manganous Oxide, Zinc Oxide, Copper Sulfate, Solium Selenite, Manganese Sulfate, Calcium Iodate, Cobalt Carbonate and Mineral Oil

Active Drug Ingredient:

Decoquinatate 27.2 g/ton

Guaranteed Analysis:

Crude Protein (min)..... 16.00%
Crude Fat (min)..... 3.00%
Crude Fiber (max)..... 15.00%

NET WT. 50 LBS. (22.67 KG)



Visit your local Alabama Farmers Cooperative to learn more or visit afcfeed.com.