



FEED

CPC Developer 14% R



Textured Beef Cattle of Pasture Feed (Medicated)

For the prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii in growing cattle fed in confinement for slaughter.

Feeding Directions:

Feed to beef cattle on pasture. Feed at a rate of up to 1% of body weight per head per day. Always feed with a free choice forage source. Always provide free choice minerals, vitamins and clean water. Do not offer free choice to any animal. Never feed wet or moldy feed.

Caution:

Always provide a source of fresh clean water, and free choice minerals. When making changes in an animal's diet, allow a 7 to 10 day transition period. Provide ample feeder space for livestock. Do not feed wet, moldy, or insect infested feed.

Ingredients:

Roughage Products, CORN GLUTEN PELLETS, Processed Grain By-Products, STEAM FLAKED CORN, Calcium Carbonate, CPC PREMIX and MTB100 55#


Active Drug Ingredient:

Monensin 29 g/ton

Guaranteed Analysis:

Crude Protein (min)..... 14.00%
Crude Fat (min)..... 2.80%
Crude Fiber (max)..... 21.00%

NET WT. 50 LBS. (22.67 KG)



Visit your local Alabama Farmers Cooperative to learn more or visit afcfeed.com.